

## **WNAR Webinar**

By James Thorson, NOAA

### ***Forecasting nonlocal climate impacts for mobile species using multivariate spatio-temporal extensions to empirical orthogonal function analysis***

It is our great pleasure to announce the next lecture in the WNAR webinar series, to be given by our distinguished colleague Dr. James Thorson from NOAA.

**Time: 9:00am - 10:00am PST (12:00 PM - 1:00 PM EST), Friday November 20, 2020**

**This is a free event and registration is required:**

<https://www.eventbrite.com/e/wnar-webinar-forecasting-climate-impacts-tickets-125183835221>

**Zoom Dial-in information and webinar link will be sent 24 hours before the event.**

#### **Abstract:**

Societal responses to COVID-19 have illustrated the great public value of accurate epidemiological forecasts; climate change has a similar potential to upend commerce and necessitates accurate decadal forecasts of community impacts. In this talk, I discuss modern extensions to Empirical Orthogonal Function (EOF) analysis involving multivariate spatio-temporal models, and how it can be used to jointly analyze climate change and community ecology. EOF analysis is widely used to identify modes of variability from spatially distributed environmental measurements (e.g., the El Niño Southern Oscillation is primary mode of variability in sea surface temperatures in the Pacific Ocean), but is less common in community (or epidemiological) modelling to understand modes of community variability. I specifically argue that EOF analysis is one potential approach to study “ecological teleconnections,” wherein individual movement generates mechanistic associations (and resulting correlations) between community dynamics occurring at geographically distant locations. I begin by adapting EOF analysis for multivariate, spatially unbalanced, zero-inflated, and right-skewed data that are common in community ecology, and show that movement of bottom-associated species in the eastern Bering Sea contains the finger-print of major climate indices. I then couple an EOF of ocean physics to spatially integrated measurements of juvenile survival, and show that this “EOF regression” can identify spatial areas that are maximally correlated with time-series residuals for recruitment as well as forecasting end-of-century recruitment. I conclude by fitting an EOF end-to-end ecosystem model to ocean physics measurements and climate-linked projections, planktonic primary and secondary producers, juvenile (pelagic) and adult (demersal) fish samples, seabirds, and fishing effort. This analysis shows greater spatial overlap between predators, juvenile, and adult walleye pollock in warm years, along with decreased overlap between adult pollock and the summer fishery, or juvenile pollock and nutritious prey. These warm conditions are projected to continue under likely climate scenarios, so resulting decadal-scale projections illustrate “community reassembly” involving this

commercially important fishery. I conclude by recommending a coordinated “Grant Challenge” for habitat science to inform spatial management, which would address these nonlocal effects, as well as additional indirect and counterfactual effects that complicate habitat science and conservation.

**Speaker Bio:**

Jim Thorson is a Senior Scientist at the National Marine Fisheries Service (NOAA Fisheries), and the Program Leader of Habitat and Ecological Processes Research at the Alaska Fisheries Science Center. His research is at the interface between statistics, ecology, and fisheries management, including applied research in spatio-temporal ecosystem modelling, life-history synthesis, and stock assessment.

WNAR is the Western North American Region of [The International Biometric Society](#), an international professional and academic society promoting the development and application of statistical and mathematical theory and methods in the biosciences. Visit [www.wnar.org](http://www.wnar.org) for more details and events.